

XMM-Newton detects the beginning of the X-ray decline of SN 1995N

P. Mucciarelli

*INAF-Astronomical Observatory of Padova
Department of Astronomy, University of Padova*

L. Zampieri

INAF-Astronomical Observatory of Padova

A. Pastorello

*INAF-Astronomical Observatory of Arcetri
Max Planck Institut für Astrophysik, Garching*

Abstract. We present the results of a new *XMM-Newton* observation of the interacting supernova 1995N, performed on July 27, 2003. We find that the 0.2-10.0 keV flux has dropped at a level of $1.44 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$, about one order of magnitude lower than that of a previous *ASCA* observation performed on January 1998. The X-ray spectral analysis shows statistically significant evidence for the presence of two distinct components, that can be modeled with emission from optically thin, thermal plasmas at different temperatures. From these temperatures we derive that the exponent of the ejecta density distribution is $n \sim 6.5$.

SN 1995N, discovered on May 1995 several months after the explosion (Benetti et al. 1995), is of special interest in the context of circumstellar medium (CSM) interacting supernovae. It is hosted in the (IB(s)m pec) galaxy MCG-02-38-017, at a distance of ~ 24 Mpc. The epoch of explosion is not known, but was estimated to be about 10 months before its optical discovery (Benetti et al. 1995).

1. *XMM-Newton* observation of SN 1995N

We observed SN 1995N with *XMM-Newton* for 72 ks on July 27, 2003, nine years after the explosion. Here we present a preliminary analysis of this observation (for details see Zampieri et al. 2004, hereafter Z04). The X-ray pointing was coordinated with near-IR and optical observations (see Pastorello et al. 2004). The X-ray data set was heavily affected by solar flares. We analysed the longer *XMM* EPIC exposures of both the MOS and pn detectors (56 and 64 ks respectively). Data were filtered using the count rate criterion $< 0.5 \text{ counts s}^{-1}$ for the EPIC MOS and $< 1.0 \text{ counts s}^{-1}$ for the EPIC pn, leaving 22 and 14 ks of useful data, respectively. Source counts were extracted from a circular region of $20''$ centered on the radio position of the supernova (van Dyk et al. 1996). Background counts were extracted from a circular region of $40''$, on the same CCD. The source count rate was $9.7 \times 10^{-3} \text{ counts s}^{-1}$ for the EPIC MOS (~ 450 counts) and $3.3 \times 10^{-2} \text{ counts s}^{-1}$ for the EPIC pn (~ 500 counts). EPIC MOS

and pn spectra were binned requiring at least 15 counts per bin. Joint MOS and pn spectral fits were performed in the 0.2-10.0 keV interval with an overall normalization constant. Despite the low statistics, the fit with single component models was not satisfactory (see Z04). The improvement obtained adding a MEKAL¹ component to a single component spectral models was significant at the $\sim 4\sigma$ level. The best fit was obtained with an absorbed double MEKAL model with column density of the interstellar medium $N_H = 1.3 \times 10^{21} \text{ cm}^{-2}$ and temperatures of the two thermal components $kT \simeq 0.8$ and 9.5 keV ($\chi^2_{red} \simeq 0.76$; see Figure 1, *left panel*).




Figure 1. *Left panel:* XMM EPIC MOS and pn spectra of SN 1995N in the 0.2-10.0 keV interval, along with the best fitting continuum model (solid line) and the two MEKAL components (dotted lines). *Right panel:* 0.2-10.0 keV unabsorbed lightcurve of SN 1995N. Fluxes were de-absorbed and, when necessary, extrapolated in the 0.2-10.0 keV interval. An estimate of the errors, based on counting statistics, is $\sim 10\%$.

We also report the detection of a faint object inside the X-ray error box of SN 1995N in the summed XMM OM UVW1 image. The object was at the limit of detectability in each single frame. Its position in the summed image is within $0.4''$ from the radio position of SN 1995N. Within the errors of the astrometric calibration, we then identify this object with SN 1995N. The lack of reference stars in the UVW1 band prevented us from performing a photometric calibration of the image. Taking the detection limit count rate of a single image as a lower bound for the UVW1 flux of SN 1995N, we estimate $F_{UVW1} \gtrsim 6.3 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$.

2. X-ray variability

Figure 1 (*right panel*) shows the fluxes derived from all the available X-ray observations of SN 1995N. The first X-ray observation was performed with ROSAT HRI on July 23, 1996 (1.3 ks; Lewin, Zimmermann & Aschenbach 1996), fol-

¹The MEKAL model is the spectrum emitted by an optically thin, thermal plasma.

lowed by other two exposures taken on August 12, 1996 (17 ks) and on August 17, 1997 (19 ks). The *ROSAT* fluxes were derived from the count rates reported by Fox et al. (2000), assuming a POWER-LAW spectrum with photon index and column density from their fit of the *ASCA* spectra. The *ASCA* observation was performed on January 19, 1998 (83 and 96 ks, respectively for the SIS and GIS instruments; Fox et al. 2000). For *ASCA* and *XMM*, the fluxes reported in Figure 1 are the average between the different instruments. It is worth noting the large decrease in the X-ray flux between the *ASCA* and *XMM* observations (unabsorbed [0.2-10.0 keV] fluxes of 1.65×10^{-12} and 1.75×10^{-13} erg cm $^{-2}$ s $^{-1}$, respectively).

3. Discussion

SN 1995N is one of the few supernovae detected in X-rays at an age of ~ 9 years. Our *XMM* observation shows that the unabsorbed X-ray flux of SN 1995N dropped at a value of 1.75×10^{-13} erg cm $^{-2}$ s $^{-1}$, about an order of magnitude lower than that of the previous *ASCA* observation performed ~ 6 years before. The decline of the X-ray flux is signalling that SN 1995N probably started to evolve towards the remnant stage. Interpreting the evolution of the X-ray light curve is not straightforward. A complex scenario in which a two-phase (clumpy and smooth) CSM contributes to the observed X-ray emission is consistent with the available data (see Z04).

The EPIC spectrum of SN 1995N shows statistically significant evidence for the presence of two distinct thermal (MEKAL) components at temperatures of 0.8 and 9.5 keV, respectively. In the standard model of ejecta/wind interaction (Chevalier & Fransson 1994), these represent the temperatures of the gas between the contact discontinuity and the reverse/forward shock. The temperature of the hotter phase is similar to the temperature of the single-component spectral fit of the *ASCA* data performed by Fox et al. (2000). Within the assumptions of the standard model, we can derive the exponent of the ejecta density distribution n from the expression $T_1/T_2 = (3-s)^2/(n-3)^2$, where s is the exponent of the CMS density distribution (Fransson, Lundqvist & Chevalier 1996). Assuming a constant and homogeneous stellar wind ($s = 2$), from the values of the temperatures T_1 and T_2 inferred from the X-ray spectrum, we obtain $n \sim 6.5$.

References

- Benetti, S., Bouchet, P., Schwarz, H., 1995, IAU Circ. 6170
- Chevalier, R.A., Fransson, C., 1994, ApJ, 420, 268
- Fox, D.W., Lewin, W.H.G., Fabian, A., et al., 2000, MNRAS, 319, 1154
- Fransson, C., Lundqvist, P., Chevalier, R.A., 1996, ApJ, 461, 993
- Lewin, W.H.G., Zimmermann, H.U., Aschenbach, B., 1996, IAU Circ. 6506
- Pastorello, A., Aretxaga, I., Zampieri, L., et al. 2004, this proceedings
- van Dyk, S.D., Sramek, R.A., Weiler, K.W., et al., 1996, IAU Circ. 6386
- Zampieri, L., Mucciarelli, P., Pastorello, A., et al., 2004, MNRAS submitted